




424-11-2019



Ranolazine Extended-Release Tablets

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use RANOLAZINE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for RANOLAZINE EXTENDED-RELEASE TABLETS.

RANOLAZINE extended-release tablets, for oral use.

Initial U.S. Approval: 2006

**INDICATIONS AND USAGE**

Ranolazine extended-release tablets are an antianginal indicated for the treatment of chronic angina. (1)

**DOSE AND ADMINISTRATION**

500 mg twice daily and increase to 1000 mg twice daily, based on clinical symptoms (2.1)

**DOSE FORMS AND STRENGTHS**

Extended-release tablets: 500 mg, 1000 mg (3)

**CONTRAINDICATIONS**

- Strong CYP3A inhibitors (e.g., ketoconazole, clarithromycin, neflavir) (4, 7.1)
- CYP3A inducers (e.g., rifampin, phenobarbital, St. John's wort) (4, 7.1)
- Liver cirrhosis (4, 8.6)

**WARNINGS AND PRECAUTIONS**

- QT interval prolongation: Can occur with ranolazine. Little data available on high doses, long exposure, use with QT interval-prolonging drugs, potassium channel variants causing prolonged QT interval, in patients with a family history of (or congenital) long QT syndrome, or in patients with known acquired QT interval prolongation (5.1)
- Renal failure: Monitor renal function after initiation and periodically in patients with moderate to severe renal impairment (CrCL<60 mL/min). If acute renal failure develops, discontinue Ranolazine extended-release tablets. (5.2)

**FULL PRESCRIBING INFORMATION: CONTENTS\*****1 INDICATIONS AND USAGE****2 DOSAGE AND ADMINISTRATION****2.1 Dosing Information****2.2 Dose Modification****3 DOSAGE FORMS AND STRENGTHS****4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS****5.1 QT Interval Prolongation****5.2 Renal Failure****6 ADVERSE REACTIONS****6.1 Clinical Trial Experience****6.2 Postmarketing Experience****7 DRUG INTERACTIONS****7.1 Effects of Other Drugs on Ranolazine****7.2 Effects of Ranolazine on Other Drugs****8 USE IN SPECIFIC POPULATIONS****8.1 Pregnancy****8.2 Lactation****8.4 Pediatric Use****8.5 Geriatric Use****ADVERSE REACTIONS**

Most common adverse reactions (>4% and more common than with placebo) are dizziness, headache, constipation, nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact ScieGen at (855) 724-3436 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

**DRUG INTERACTIONS**

- Moderate CYP3A inhibitors (e.g., diltiazem, verapamil, erythromycin): Limit Ranolazine extended-release tablets to 500 mg twice daily. (7.1)
- P-gp inhibitors (e.g., cyclosporine): Ranolazine exposure increased. Titrate Ranolazine extended-release tablets based on clinical response. (7.1)
- CYP3A substrates: Limit simvastatin to 20 mg when used with Ranolazine extended-release tablets. Doses of other sensitive CYP3A substrates (e.g., lovastatin) and CYP3A substrates with narrow therapeutic range (e.g., cyclosporine, tacrolimus, sirolimus) may need to be reduced with Ranolazine extended-release tablets. (7.2)
- OCT2 substrates: Limit the dose of metformin to 1700 mg daily when used with Ranolazine extended-release tablets 1000 mg twice daily. Doses of other OCT2 substrates may require adjusted doses. (7.2)
- Drugs transported by P-gp (e.g., digoxin), or drugs metabolized by CYP2D6 (e.g., tricyclic antidepressants) may need reduced doses when used with Ranolazine extended-release tablets. (7.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-Approved Patient Labeling.

Revised: 11/2019

8.6 Use in Patients with Hepatic Impairment

8.7 Use in Patients with Renal Impairment

8.8 Use in Patients with Heart Failure

8.9 Use in Patients with Diabetes Mellitus

**10 OVERDOSAGE****11 DESCRIPTION****12 CLINICAL PHARMACOLOGY****12.1 Mechanism of Action****12.2 Pharmacodynamics****12.3 Pharmacokinetics****13 NONCLINICAL TOXICOLOGY****13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility****14 CLINICAL STUDIES****14.1 Chronic Stable Angina****14.2 Lack of Benefit in Acute Coronary Syndrome****15 REFERENCES****16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

**FULL PRESCRIBING INFORMATION****1 INDICATIONS AND USAGE**

Ranolazine extended-release tablets are indicated for the treatment of chronic angina.

Ranolazine extended-release tablets may be used with beta-blockers, nitrates, calcium channel blockers, anti-platelet therapy, lipid-lowering therapy, ACE inhibitors, and angiotensin receptor blockers.

**2 DOSAGE AND ADMINISTRATION****2.1 Dosing Information**

Initiate Ranolazine extended-release tablets dosing at 500 mg twice daily and increase to 1000 mg twice daily, as needed, based on clinical symptoms. Take Ranolazine extended-release tablets with or without meals. Swallow Ranolazine extended-release tablets whole; do not crush, break, or chew.

The maximum recommended daily dose of Ranolazine extended-release tablets is 1000 mg twice daily. If a dose of Ranolazine extended-release tablets is missed, take the prescribed dose at the next scheduled time; do not double the next dose.

**2.2 Dose Modification**

Dose adjustments may be needed when Ranolazine extended-release tablets are taken in combination with certain other drugs [see Drug Interactions (7.1)]. Limit the maximum dose of Ranolazine extended-release tablets to 500 mg twice daily in patients on moderate CYP3A inhibitors such as diltiazem, verapamil, and erythromycin. Use of Ranolazine extended-release tablets with strong CYP3A inhibitors is contraindicated [see Contraindications (4), Drug Interactions (7.1)]. Use of P-gp inhibitors, such as cyclosporine, may increase exposure to Ranolazine extended-release tablets. Titrate Ranolazine extended-release tablets based on clinical response [see Drug Interactions (7.1)].

**3 DOSAGE FORMS AND STRENGTHS**

Ranolazine extended-release tablets are in the following strengths:

- Peach, oblong shaped, film-coated, extended release tablet debossed with "423" on one side and "SG" on the other side.
- Pale yellow, oblong shaped, film-coated, extended release tablet debossed with "424" on one side and "SG" on the other side.

**4 CONTRAINDICATIONS**

Ranolazine extended-release tablets are contraindicated in patients:

- Taking strong inhibitors of CYP3A [see Drug Interactions (7.1)]
- Taking inducers of CYP3A [see Drug Interactions (7.1)]
- With liver cirrhosis [see Use in Specific Populations (8.6)]

**5 WARNINGS AND PRECAUTIONS****5.1 QT Interval Prolongation**

Ranolazine blocks  $I_{Kr}$  and prolongs the QTc interval in a dose-related manner. Clinical experience in an acute coronary syndrome population did not show an increased risk of proarrhythmia or sudden death [see Clinical Studies (14.2)]. However, there is little experience with high doses (>1000 mg twice daily) or exposure, other QT-prolonging drugs, potassium channel variants resulting in a long QT interval, in patients with a family history of (or congenital) long QT syndrome, or in patients with known acquired QT interval prolongation.

**5.2 Renal Failure**

Acute renal failure has been observed in some patients with severe renal impairment (creatinine clearance [CrCL] <30 mL/min) while taking Ranolazine extended-release tablets. If acute renal failure develops (e.g., marked increase in serum creatinine associated with an increase in blood urea nitrogen [BUN]), discontinue Ranolazine extended-release tablets and treat appropriately [see Use in Specific Populations (8.7)].

**6 ADVERSE REACTIONS****6.1 Clinical Trial Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

A total of 2018 patients with chronic angina were treated with ranolazine in controlled clinical trials. Of the patients treated with Ranolazine extended-release tablets, 1026 were enrolled in three double-blind, placebo-controlled, randomized studies (CARISA, ERICA, MARISA) of up to 12 weeks' duration. In addition, upon study completion, 1251 patients received treatment with Ranolazine extended-release tablets in open-label, long-term studies; 1227 patients were exposed to Ranolazine extended-release tablets for more than 1 year, 613 patients for more than 2 years, 531 patients for more than 3 years, and 326 patients for more than 4 years.

At recommended doses, about 6% of patients discontinued treatment with Ranolazine extended-release tablets because of an adverse event in controlled studies in angina patients compared to about 3% on placebo. The most common adverse events that led to discontinuation more frequently on Ranolazine extended-release tablets than placebo were dizziness (1.3% versus 0.1%), nausea (1% versus 0%), asthenia, constipation, and headache (each about 0.5% versus 0%). Doses above 1000 mg twice daily are poorly tolerated.

In controlled clinical trials of angina patients, the most frequently reported treatment-emergent adverse reactions (>4% and more common on Ranolazine extended-release tablets than on placebo) were dizziness (6.2%), headache (5.5%), constipation (4.5%), and nausea (4.4%). Dizziness may be dose-related. In open-label, long-term treatment studies, a similar adverse reaction profile was observed.

The following additional adverse reactions occurred at an incidence of 0.5 to 4.0% in patients treated with Ranolazine extended-release tablets and were more frequent than the incidence observed in placebo-treated patients:

**Cardiac Disorders** – bradycardia, palpitations

**Ear and Labyrinth Disorders** – tinnitus, vertigo

**Eye Disorders** – blurred vision

**Gastrointestinal Disorders** – abdominal pain, dry mouth, vomiting, dyspepsia

**General Disorders and Administrative Site Adverse Events** – asthenia, peripheral edema

**Metabolism and Nutrition Disorders** – anorexia

**Nervous System Disorders** – syncope (vasovagal)

**Psychiatric Disorders** – confusional state

**Renal and Urinary Disorders** – hematuria

**Respiratory, Thoracic, and Mediastinal Disorders** – dyspnea

**Skin and Subcutaneous Tissue Disorders** – hyperhidrosis

**Vascular Disorders** – hypotension, orthostatic hypotension

Other (<0.5%) but potentially medically important adverse reactions observed more frequently with Ranolazine extended-release tablets than placebo treatment in all controlled studies included: angioedema, renal failure, eosinophilia, chromaturia, blood urea increased, hypoesthesia, paresthesia, tremor, pulmonary fibrosis, thrombocytopenia, leukopenia, and pancytopenia.

A large clinical trial in acute coronary syndrome patients was unsuccessful in demonstrating a benefit for Ranolazine extended-release tablets, but there was no apparent proarrhythmic effect in these high-risk patients [see Clinical Studies (14.2)].

**Laboratory Abnormalities:**

Ranolazine extended-release tablets produces elevations of serum creatinine by 0.1 mg/dL, regardless of previous renal function, likely because of inhibition of creatinine's tubular secretion. In general, the elevation has a rapid onset, shows no signs of progression during long-term therapy, is reversible after discontinuation of Ranolazine extended-release tablets, and is not accompanied by changes in BUN. In healthy volunteers, Ranolazine extended-release tablets 1000 mg twice daily had no effect upon the glomerular filtration rate. More marked and progressive increases in serum creatinine, associated with increases in BUN or potassium, indicating acute renal failure, have been reported after initiation of Ranolazine extended-release tablets in patients with severe renal impairment [see Warnings and Precautions (5.2), Use in Specific Populations (8.7)].

**6.2 Postmarketing Experience**

The following adverse reactions have been identified during postapproval use of Ranolazine extended-release tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure:

**Nervous System Disorders** – Abnormal coordination, myoclonus, paresthesia, tremor, and other serious neurologic adverse events have been reported to occur, sometimes concurrently, in patients taking ranolazine. The onset of events was often associated with an increase in ranolazine dose or exposure. Many patients reported symptom resolution following drug discontinuation or dose decrease.

**Metabolism and Nutrition Disorders** – Cases of hypoglycemia have been reported in diabetic patients on anti-diabetic medication.

**Psychiatric Disorders** – hallucination

**Renal and Urinary Disorders** – dysuria, urinary retention

**Skin and Subcutaneous Tissue Disorders** – angioedema, pruritus, rash

**7 DRUG INTERACTIONS****7.1 Effects of Other Drugs on Ranolazine**

**Strong CYP3A Inhibitors**  
Do not use Ranolazine extended-release tablets with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, neflavir, ritonavir, indinavir, and saquinavir [see Contraindications (4), Clinical Pharmacology (12.3)].

**Moderate CYP3A Inhibitors**

Limit the dose of Ranolazine extended-release tablets to 500 mg twice daily in patients on moderate CYP3A inhibitors, including diltiazem, verapamil, erythromycin, fluconazole, and grapefruit juice or grapefruit-containing products [see Dosage and Administration (2.2), Clinical Pharmacology (12.3)].

**P-gp Inhibitors**

Concomitant use of Ranolazine extended-release tablets and P-gp inhibitors, such as cyclosporine, may result in increases in ranolazine concentrations. Titrate Ranolazine extended-release tablets based on clinical response in patients concomitantly treated with predominant P-gp inhibitors such as cyclosporine [see Dosage and Administration (2.2)].

**CYP3A Inducers**

Do not use Ranolazine extended-release tablets with CYP3A inducers such as rifampin, rifabutin, rifapentine, phenobarbital, phenytoin, carbamazepine, and St. John's wort [see Contraindications (4), Clinical Pharmacology (12.3)].

**7.2 Effects of Ranolazine on Other Drugs****Drugs Metabolized by CYP3A**

Limit the dose of simvastatin in patients on any dose of Ranolazine extended-release tablets to 20 mg once daily, when ranolazine is co-administered. Dose adjustment of other sensitive CYP3A substrates (e.g., lovastatin) and CYP3A substrates with a narrow therapeutic range (e.g., cyclosporine, tacrolimus, sirolimus) may be required as Ranolazine extended-release tablets may increase plasma concentrations of these drugs [see Clinical Pharmacology (12.3)].

**Drugs Transported by P-gp**

Concomitant use of ranolazine and digoxin results in increased exposure to digoxin. The dose of digoxin may have to be adjusted [see Clinical Pharmacology (12.3)].

**Drugs Metabolized by CYP2D6**

The exposure to CYP2D6 substrates, such as tricyclic antidepressants and antipsychotics, may be increased during co-administration with Ranolazine extended-release tablets, and lower doses of these drugs may be required.

**Drugs Transported by OCT2**

In subjects with type 2 diabetes mellitus, concomitant use of Ranolazine extended-release tablets 1000 mg twice daily and metformin results in increased plasma levels of metformin. When Ranolazine extended-release tablets 1000 mg twice daily is co-administered with metformin, metformin dose should not exceed 1700 mg/day. Monitor blood glucose levels and risks associated with high exposures of metformin.

Metformin exposure was not significantly increased when given with Ranolazine extended-release tablets 500 mg twice daily [see Clinical Pharmacology (12.3)].

**8 USE IN SPECIFIC POPULATIONS****8.1 Pregnancy****Risk Summary**

There are no available data on Ranolazine extended-release tablets use in pregnant women to inform any drug-associated risks. Studies in rats and rabbits showed no evidence of fetal harm at exposures 4 times the maximum recommended human dose (MRHD) [see Data].

In the U.S. general population, the estimated background risk of major birth defects and of miscarriage of clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

**Data****Animal Data**

Embryofetal toxicity studies were conducted in rats and rabbits orally administered ranolazine during organogenesis. In rats, decreased fetal weight and reduced ossification were observed at doses (corresponding to 4-fold the AUC for the MRHD) that caused maternal weight loss. No adverse fetal effects were observed in either species exposed (AUC) to ranolazine at exposures (AUC) equal to the MRHD.

**8.2 Lactation****Risk Summary**

There are no data on the presence of ranolazine in human milk, the effects on the breastfed infant, or the effects on milk production. However, ranolazine is present in rat milk [see Use in Specific Populations (8.1)]. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Ranolazine extended-release tablets and any potential adverse effects on the breastfed infant from Ranolazine extended-release tablets or from the underlying maternal condition.

Adult female rats were administered ranolazine orally from gestation day 6 through postnatal day 20. No adverse effects on pup development, behavior, or reproduction parameters were observed at a maternal dosage level of 60 mg/kg/day (equal to the MRHD based on AUC). At maternally toxic doses, male and female pups exhibited increased mortality and decreased body weight, and female pups showed increased motor activity. The pups were potentially exposed to low amounts of ranolazine via the maternal milk.

**8.4 Pediatric Use**

Safety and effectiveness have not been established in pediatric patients.

**8.5 Geriatric Use**

Of the chronic angina patients treated with Ranolazine extended-release tablets in controlled studies, 496 (48%) were ≥65 years of age, and 114 (11%) were ≥75 years of age. No overall differences in efficacy were observed between older and younger patients. There were no differences in safety for patients ≥65 years compared to younger patients, but patients ≥75 years of age on Ranolazine extended-release tablets, compared to placebo, had a higher incidence of adverse events, serious adverse events, and drug discontinuations due to adverse events. In general, dose selection for an elderly patient should usually start at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease, or other drug therapy.

**8.6 Use in Patients with Hepatic Impairment**

Ranolazine extended-release tablets are contraindicated in patients with liver cirrhosis. In a study of cirrhotic patients, the  $C_{max}$  of ranolazine was increased 30% in cirrhotic patients with mild (Child-Pugh Class A) hepatic impairment, but increased 80% in cirrhotic patients with moderate (Child-Pugh Class B) hepatic impairment compared to patients without hepatic impairment. This increase was not enough to account for the 3-fold increase in QT prolongation seen in cirrhotic patients with mild to moderate hepatic impairment [see Clinical Pharmacology (12.2)].

**8.7 Use in Patients with Renal Impairment**

A pharmacokinetic study of Ranolazine extended-release tablets in subjects with severe renal impairment (CrCL <30 mL/min) was stopped when 2 of 4 subjects developed acute renal failure after receiving Ranolazine extended-release tablets 500 mg twice daily for 5 days (lead-in phase) followed by 1000 mg twice a day (1 dose in one subject and 11 doses in the other). Increases in creatinine, BUN, and potassium were observed in 3 subjects during the 500 mg lead-in phase. One subject required hemodialysis, while the other 2 subjects improved upon drug discontinuation [see Warnings and Precautions (5.2)]. Monitor renal function periodically in patients with moderate to severe renal impairment. Discontinue Ranolazine extended-release tablets if acute renal failure develops.

In a separate study,  $C_{max}$  was increased between 40% and 50% in patients with mild, moderate, or severe renal impairment compared to patients with no renal impairment, suggesting a similar increase in exposure in patients with renal failure independent of the degree of impairment. The pharmacokinetics of ranolazine has not been assessed in patients on dialysis.

**8.8 Use in Patients with Heart Failure**

Heart failure (NYHA Class I to IV) had no significant effect on ranolazine pharmacokinetics. Ranolazine extended-release tablets had minimal effects on heart rate and blood pressure in patients with angina and heart failure NYHA Class I to IV. No dose adjustment of Ranolazine extended-release tablets is required in patients with heart failure.

**8.9 Use in Patients with Diabetes Mellitus**

A population pharmacokinetic evaluation of data from angina patients and healthy subjects showed no effect of diabetes on ranolazine pharmacokinetics. No dose adjustment is required in patients with diabetes.

Ranolazine extended-release tablets produces small reductions in HbA1c in patients with diabetes, the clinical significance of which is unknown. Ranolazine extended-release tablets should not be considered a treatment for diabetes.


**10 OVERDOSAGE**

Hypotension, QT prolongation, bradycardia, myoclonic activity, severe tremor, unsteady gait/incoordination, dizziness, nausea, vomiting, dysphasia, and hallucinations have been seen in cases of oral overdose of Ranolazine extended-release tablets. In cases of extreme overdose of Ranolazine extended-release tablets fatal outcomes have been reported. In clinical studies, high intravenous exposure resulted in diopnea, paresthesia, confusion, and syncope. In addition to general supportive measures, continuous ECG monitoring may be warranted in the event of overdose. Since Ranolazine is about 62% bound to plasma proteins, hemodialysis is unlikely to be effective in clearing Ranolazine.

**11 DESCRIPTION**

Ranolazine is available as a film-coated, extended-release tablet for oral administration.

Ranolazine is a racemic mixture, chemically described as 1-piperazineacetamide, *N*-(2,6-dimethylphenyl)-4-[2-hydroxy-3-(2-methoxyphenoxy)propyl]-, (±). It has an empirical formula of  $C_{24}H_{33}N_3O_4$ , a molecular weight of 427.54 g/mole, and the following structural formula:



Ranolazine is a white to off-white crystalline powder. Ranolazine is soluble in dichloromethane and methanol.

Ranolazine extended-release tablets contain 500 mg or 1000 mg of ranolazine and the following inactive ingredients: microcrystalline cellulose, methacrylic acid and ethyl acrylate copolymer, sodium lauryl sulfate, polysorbate 80, hypromellose, sodium hydroxide and magnesium stearate. Additional inactive ingredients for the 500 mg tablet include hydroxypropyl methyl cellulose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin, Iron Oxide Yellow, and Iron Oxide Red; additional inactive ingredients for the 1000 mg tablet include hydroxypropyl methyl cellulose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin and Iron Oxide Yellow.

**12 CLINICAL PHARMACOLOGY****12.1 Mechanism of Action**

The mechanism of action of ranolazine's antianginal effects has not been determined. Ranolazine has anti-ischemic and antianginal effects that do not depend upon reductions in heart rate or blood pressure. It does not affect the rate-pressure product, a measure of myocardial work, at maximal exercise. Ranolazine at therapeutic levels can inhibit the cardiac late sodium current ( $I_{NaL}$ ). However, the relationship of this inhibition to angina symptoms is uncertain.


The QT prolongation effect of ranolazine on the surface electrocardiogram is the result of inhibition of  $I_{Kr}$ , which prolongs the ventricular action potential.

**12.2 Pharmacodynamics****Hemodynamic Effects**

Patients with chronic angina treated with Ranolazine extended-release tablets in controlled clinical studies had minimal changes in mean heart rate (<2 bpm) and systolic blood pressure (<3 mm Hg). Similar results were observed in subgroups of patients with CHF NYHA Class I or II, diabetes, or reactive airway disease, and in elderly patients.

**Electrocardiographic Effects**

Dose and plasma concentration-related increases in the QTc interval [see Warnings and Precautions (5.1)], reductions in T wave amplitude, and, in some cases, notched T waves, have been observed in patients treated with Ranolazine extended-release tablets. These effects are believed to be caused by ranolazine and not by its metabolites. The relationship between the change in QTc and ranolazine plasma concentrations is linear, with a slope of about 2.6 msec/1000 ng/mL, through exposures corresponding to doses several-fold higher than the maximum recommended dose of 1000 mg twice daily. The variable blood levels attained after a given dose of ranolazine give a wide range of effects on QTc. At  $T_{max}$  following repeat dosing at 1000 mg twice daily, the mean change in QTc is about 6 msec



### OCT2 Substrates

In subjects with type 2 diabetes mellitus, the exposure to metformin is increased by 40% and 80% following administration of ranolazine 500 mg twice daily and 1000 mg twice daily, respectively. If co-administered with Ranolazine extended-release tablets 1000 mg twice daily, do not exceed metformin doses of 1700 mg/day [see Drug Interactions (7.2)].

### 13 NONCLINICAL TOXICOLOGY

#### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Ranolazine tested negative for genotoxic potential in the following assays: Ames bacterial mutation assay, Saccharomyces assay for mitotic gene conversion, chromosomal aberrations assay in Chinese hamster ovary (CHO) cells, mammalian CHO/HGPRT gene mutation assay, and mouse and rat bone marrow micronucleus assays.

There was no evidence of carcinogenic potential in mice or rats. The highest oral doses used in the carcinogenicity studies were 150 mg/kg/day for 21 months in rats (900 mg/m<sup>2</sup>/day) and 50 mg/kg/day for 24 months in mice (150 mg/m<sup>2</sup>/day). These maximally tolerated doses are 0.8 and 0.1 times, respectively, the daily maximum recommended human dose (MRHD) of 2000 mg on a surface area basis. A published study reported that ranolazine promoted tumor formation and progression to malignancy when given to transgenic APC (min/+ ) mice at a dose of 30 mg/kg twice daily [see References (15)]. The clinical significance of this finding is unclear.

In male and female rats, oral administration of ranolazine that produced exposures (AUC) approximately 3-fold or 5-fold higher, respectively, than the MRHD had no effect on fertility.

### 14 CLINICAL STUDIES

#### 14.1 Chronic Stable Angina

CARISA (Combination Assessment of Ranolazine In Stable Angina) was a study in 823 chronic angina patients randomized to receive 12 weeks of treatment with twice-daily Ranolazine extended-release tablets 750 mg, 1000 mg, or placebo, who also continued daily doses of atenolol 50 mg, amlodipine 5 mg, or diltiazem CD 180 mg. Sublingual nitrates were used in this study as needed.

In this trial, statistically significant (p < 0.05) increases in modified Bruce treadmill exercise duration and time to angina were observed for each Ranolazine extended-release tablets dose versus placebo, at both trough (12 hours after dosing) and peak (4 hours after dosing) plasma levels, with minimal effects on blood pressure and heart rate. The changes versus placebo in exercise parameters are presented in Table 1. Exercise treadmill results showed no increase in effect on exercise at the 1000 mg dose compared to the 750 mg dose.

Table 1 Exercise Treadmill Results (CARISA)

Study	Mean Difference from Placebo (sec)		
	750 mg	1000 mg	
Exercise Duration			
	Trough	24*	24*
	Peak	34*	26*
Time to Angina			
	Trough	30*	26*
	Peak	38 <sup>b</sup>	38 <sup>b</sup>
Time to 1 mm ST-Segment Depression			
	Trough	20	21
	Peak	41 <sup>b</sup>	35 <sup>b</sup>

\* p-value ≤ 0.05      <sup>b</sup> p-value ≤ 0.005

The effects of Ranolazine extended-release tablets on angina frequency and nitroglycerin use are shown in Table 2.

Table 2 Angina Frequency and Nitroglycerin Use (CARISA)

	N	Placebo	Ranolazine extended-release tablets 750 mg <sup>a</sup>	Ranolazine extended-release tablets 1000 mg <sup>a</sup>
Angina Frequency (attacks/week)	N	258	272	261
	Mean	3.3	2.5	2.1
	P-value vs Placebo	—	0.006	< 0.001
Nitroglycerin Use (doses/week)	N	252	262	244
	Mean	3.1	2.1	1.8
	P-value vs placebo	—	0.016	< 0.001

<sup>a</sup> Twice daily

Tolerance to Ranolazine extended-release tablets did not develop after 12 weeks of therapy. Rebound increases in angina, as measured by exercise duration, have not been observed following abrupt discontinuation of Ranolazine extended-release tablets.

Ranolazine extended-release tablets has been evaluated in patients with chronic angina who remained symptomatic despite treatment with the maximum dose of an antianginal agent. In the ERICA (Efficacy of Ranolazine In Chronic Angina) trial, 565 patients were randomized to receive an initial dose of Ranolazine extended-release tablets 500 mg twice daily or placebo for 1 week, followed by 6 weeks of treatment with Ranolazine extended-release tablets 1000 mg twice daily or placebo, in addition to concomitant treatment with amlodipine 10 mg once daily. In addition, 45% of the study population also received long-acting nitrates. Sublingual nitrates were used as needed to treat angina episodes. Results are shown in Table 3. Statistically significant decreases in angina attack frequency (p=0.028) and nitroglycerin use (p=0.014) were observed with Ranolazine extended-release tablets compared to placebo. These treatment effects appeared consistent across age and use of long-acting nitrates.

Table 3 Angina Frequency and Nitroglycerin Use (ERICA)

	N	Placebo	Ranolazine extended-release tablets <sup>a</sup>
Angina Frequency (attacks/week)	N	281	277
	Mean	4.3	3.3
	Median	2.4	2.2
Nitroglycerin Use (doses/week)	N	281	277
	Mean	3.6	2.7
	Median	1.7	1.3

<sup>a</sup> 1000 mg twice daily

### Gender

Effects on angina frequency and exercise tolerance were considerably smaller in women than in men. In CARISA, the improvement in Exercise Tolerance Test (ETT) in females was about 33% of that in males at the 1000 mg twice-daily dose level. In ERICA, where the primary endpoint was angina attack frequency, the mean reduction in weekly angina attacks was 0.3 for females and 1.3 for males.

### Race

There were insufficient numbers of non-Caucasian patients to allow for analyses of efficacy or safety by racial subgroup.

### 14.2 Lack of Benefit in Acute Coronary Syndrome

In a large (n=6560) placebo-controlled trial (MERLIN-TIMI 36) in patients with acute coronary syndrome, there was no benefit shown on outcome measures. However, the study is somewhat reassuring regarding proarrhythmic risks, as ventricular arrhythmias were less common on ranolazine [see Clinical Pharmacology (12.2)], and there was no difference between Ranolazine extended-release tablets and placebo in the risk of all-cause mortality (relative risk ranolazine: placebo 0.99 with an upper 95% confidence limit of 1.22).

### 15 REFERENCES

M.A. Suckow et al. The anti-ischemia agent ranolazine promotes the development of intestinal tumors in APC (min/+ ) mice. Cancer Letters 209(2004):165-9.

### 16 HOW SUPPLIED/STORAGE AND HANDLING

Ranolazine extended-release tablets are supplied as film-coated, oblong-shaped, extended-release tablets in the following strengths:

- 500 mg tablets are peach, with "423" on one side and "SG" on the other side
- 1000 mg tablets are pale yellow, with "424" on one side and "SG" on the other side

Ranolazine extended-release tablets 500 mg are available in:

Bottle of 30 Tablets: NDC 50228-423-30

Bottle of 60 Tablets: NDC 50228-423-60

Bottle of 500 Tablets: NDC 50228-423-05

Ranolazine extended-release tablets 1000 mg are available in:

Bottle of 30 Tablets: NDC 50228-424-30

Bottle of 60 Tablets: NDC 50228-424-60

Bottle of 500 Tablets: NDC 50228-424-05

Store at 25°C (77°F); with excursions permitted to 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature.]

### 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Inform patients that Ranolazine extended-release tablets will not abate an acute angina episode.

#### Strong CYP3A Inhibitors, CYP3A Inducers, Liver Cirrhosis

- Inform patients that Ranolazine extended-release tablets should not be used with drugs that are strong CYP3A inhibitors (e.g., ketoconazole, clarithromycin, nefazodone, ritonavir) [see Contraindications (4), Drug Interactions (7.1)].
- Inform patients that Ranolazine extended-release tablets should not be used with drugs that are inducers of CYP3A (e.g., rifampin, rifabutin, rifapentine, barbiturates, carbamazepine, phenytoin, St. John's wort) [see Contraindications (4), Drug Interactions (7.1)].
- Inform patients that Ranolazine extended-release tablets should not be used in patients with liver cirrhosis [see Contraindications (4), Use in Specific Populations (8.6)].

#### Moderate CYP3A Inhibitors, P-gp Inhibitors, Grapefruit Products

- Advise patients to inform their physician if they are receiving drugs that are moderate CYP3A inhibitors (e.g., diltiazem, verapamil, erythromycin) [see Drug Interactions (7)].
- Advise patients to inform their physician if they are receiving drugs that are P-gp inhibitors (e.g., cyclosporine) [see Drug Interactions (7)].
- Advise patients to limit grapefruit juice or grapefruit products when taking Ranolazine extended-release tablets [see Drug Interactions (7)].

#### QT Interval Prolongation

- Inform patients that Ranolazine extended-release tablets may produce changes in the electrocardiogram (QTc interval prolongation) [see Warnings and Precautions (5.1)].
- Advise patients to inform their physician of any personal or family history of QTc prolongation, congenital long QT syndrome, or if they are receiving drugs that prolong the QTc interval such as Class Ia (e.g., quinidine) or Class III (e.g., dofetilide, sotalol, amiodarone) antiarrhythmic agents, erythromycin, and certain antipsychotics (e.g., thioridazine, ziprasidone) [see Warnings and Precautions (5.1)].

### Use in Patients with Renal Impairment

Patients with severe renal impairment may be at risk of renal failure while on Ranolazine extended-release tablets. Advise patients to inform their physician if they have impaired renal function before or while taking Ranolazine extended-release tablets [see Warnings and Precautions (5.2)].

### Dizziness, Fainting

- Inform patients that Ranolazine extended-release tablets may cause dizziness and lightheadedness. Patients should know how they react to Ranolazine extended-release tablets before they operate an automobile or machinery, or engage in activities requiring mental alertness or coordination [see Adverse Reactions (6.1)].
- Advise patients to contact their physician if they experience fainting spells while taking Ranolazine extended-release tablets.

### Administration

- Instruct patients to swallow Ranolazine extended-release tablets whole, with or without meals, and not to crush, break, or chew tablets. Inform patients that if a dose is missed, to take the usual dose at the next scheduled time. The next dose should not be doubled. Inform patients that doses of Ranolazine extended-release tablets higher than 1000 mg twice daily should not be used [see Dosage and Administration (2)].
- Advise patients to inform their physician of any other medications taken concurrently with Ranolazine extended-release tablets, including over-the-counter medications

Manufactured by:  
ScieGen Pharmaceuticals Inc  
Hauppauge, NY 11788  
USA

Rev. 11/19

## Patient Information Ranolazine extended-release tablets Dosing Strengths: 500 mg tablets 1000 mg tablets

Read this Patient Information before you start taking Ranolazine extended-release tablets and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or treatment.

### What are Ranolazine extended-release tablets?

Ranolazine extended-release tablets are prescription medicine used to treat angina that keeps coming back (chronic angina).

Ranolazine extended-release tablets may be used with other medicines that are used for heart problems and blood pressure control.

It is not known if Ranolazine extended-release tablets are safe and effective in children.

### Who should not take Ranolazine extended-release tablets? Do not take Ranolazine extended-release tablets if:

- you take any of the following medicines:
  - for fungus infection: ketoconazole (Nizoral<sup>®</sup>), itraconazole (Sporanox<sup>®</sup>, Onmel<sup>™</sup>)
  - for infection: clarithromycin (Biaxin<sup>®</sup>)
  - for depression: nefazodone
  - for HIV: nelfinavir (Viracept<sup>®</sup>), ritonavir (Norvir<sup>®</sup>), lopinavir and ritonavir (Kaletra<sup>®</sup>), indinavir (Crixivan<sup>®</sup>), saquinavir (Invirase<sup>®</sup>)
  - for tuberculosis (TB): rifampin (Rifadin<sup>®</sup>), rifabutin (Mycobutin<sup>®</sup>), rifapentine (Priftin<sup>®</sup>)
  - for seizures: phenobarbital, phenytoin (Dilantin<sup>®</sup>, Dilantin<sup>®</sup>), Dilantin125<sup>®</sup>), carbamazepine (Tegretol<sup>®</sup>)
  - St. John's wort (Hypericum perforatum)
- you have scarring (cirrhosis) of your liver

### What should I tell my doctor before taking Ranolazine extended-release tablets?

Before you take Ranolazine extended-release tablets, tell your doctor if you:

- have or have a family history of a heart problem, called 'QT prolongation' or 'long QT syndrome'.
- have liver problems
- have kidney problems
- are pregnant or plan to become pregnant. It is not known if Ranolazine extended-release tablets will harm your unborn baby.
- are breast-feeding or plan to breast-feed. It is not known if Ranolazine passes into your breast milk. You and your doctor should decide if you will breast-feed.

Tell your doctor about all the medicines you take, including all prescription and nonprescription medicines, vitamins, and herbal supplements. Ranolazine extended-release tablets may affect the way other medicines and other medicines may affect how Ranolazine extended-release tablets works.

### Tell your doctor if you take medicines:

- for your heart
- for cholesterol
- for diabetes
- for infection
- for fungus
- for transplant
- for nausea and vomiting because of cancer treatments
- for mental problems

Know the medicines you take. Keep a list of them to show your doctor or pharmacist when you get a new medicine.

### How should I take Ranolazine extended-release tablets?

- Take Ranolazine extended-release tablets exactly as your doctor tells you.
- Your doctor will tell you how much Ranolazine extended-release tablets to take and when to take it.
- Do not change your dose unless your doctor tells you to.
- Tell your doctor if you still have symptoms of angina after starting Ranolazine extended-release tablets.
- Take Ranolazine extended-release tablets by mouth, with or without food.
- Swallow the Ranolazine extended-release tablets whole. Do not crush, break, or chew Ranolazine extended-release tablets before swallowing.
- If you miss a dose of Ranolazine extended-release tablets, wait to take the next dose of Ranolazine extended-release tablets at your regular time. Do not make up for the missed dose. Do not take more than 1 dose at a time.
- If you take too much Ranolazine extended-release tablets, call your doctor, or go to the nearest emergency room right away.

### What should I avoid while taking Ranolazine extended-release tablets?

- Grapefruit and grapefruit juice. Limit products that have grapefruit in them. They can cause your blood levels of Ranolazine extended-release tablets to increase.
- Ranolazine extended-release tablets can cause dizziness, lightheadedness, or fainting. If you have these symptoms, do not drive a car, use machinery, or do anything that needs you to be alert.

### What are the possible side effects of Ranolazine extended-release tablets?

Ranolazine extended-release tablets may cause serious side effects, including:

- changes in the electrical activity of your heart called QT prolongation. Your doctor may check the electrical activity of your heart with an ECG. Tell your doctor right away if you feel faint, lightheaded, or feel your heart beating irregularly or fast while taking Ranolazine extended-release tablets. These may be symptoms related to QT prolongation.
- kidney failure in people who already have severe kidney problems. Your doctor may need to do tests to check how your kidneys are working.

The most common side effects of Ranolazine extended-release tablets include:

- dizziness
- headache
- constipation
- nausea

Tell your doctor if you have any side effect that bothers you or does not go away.

These are not all the possible side effects of Ranolazine extended-release tablets. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### How should I store Ranolazine extended-release tablets?

Store Ranolazine extended-release tablets at room temperature between 59° to 86°F (15° to 30°C).

### Keep Ranolazine extended-release tablets and all medicines out of the reach of children. General information about Ranolazine extended-release tablets.

Medicines are sometimes prescribed for purposes other than those listed in the Patient Information. Do not use Ranolazine extended-release tablets for a condition for which it was not prescribed. Do not give Ranolazine extended-release tablets to other people, even if they have the same condition you have. It may harm them.

The Patient Information summarizes the most important information about Ranolazine extended-release tablets. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about Ranolazine extended-release tablets that is written for health professionals.

For more information call contact ScieGen at (855) 724-3436.

### What is chronic angina?

Chronic angina means pain or discomfort in the chest, jaw, shoulder, back, or arm that keeps coming back. There are other possible signs and symptoms of angina including shortness of breath. Angina usually comes on when you are active or under stress. Chronic angina is a symptom of a heart problem called coronary heart disease (CHD), also known as coronary artery disease (CAD). When you have CHD, the blood vessels in your heart become stiff and narrow. Oxygen-rich blood cannot reach your heart muscle easily. Angina comes on when too little oxygen reaches your heart muscle.

### What are the ingredients in Ranolazine extended-release tablets? Active ingredient: Ranolazine

#### Inactive ingredients:

**500 mg tablet:** microcrystalline cellulose, methacrylic acid and ethyl acrylate copolymer, sodium lauryl sulfate, polysorbate 80, hypromellose, sodium hydroxide, magnesium stearate, hydroxypropyl methyl cellulose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin, Iron oxide red, Iron oxide yellow.

**1000 mg tablet:** microcrystalline cellulose, methacrylic acid and ethyl acrylate copolymer, sodium lauryl sulfate, polysorbate 80, hypromellose, sodium hydroxide, magnesium stearate, hydroxypropyl methyl cellulose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin, Iron oxide yellow.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Manufactured by:  
ScieGen Pharmaceuticals Inc  
Hauppauge, NY 11788  
USA

Rev. 11/19

All trademarks are the property of their respective owners.

