

Width: 17.0"
Length: 24.0"
Fold: 1.25" x 1.25"

8.6 Geriatric Use
In controlled clinical studies of pregabalin in neuropathic pain associated with diabetic peripheral neuropathy, 246 patients were 65 to 74 years of age, and 73 patients were 75 years of age or older.

In controlled clinical studies of pregabalin in neuropathic pain associated with postherpetic neuralgia, 262 patients were 65 to 74 years of age, and 373 patients were 75 years of age or older. In controlled clinical studies of pregabalin in epilepsy, there were only 10 patients 65 to 74 years of age, and 2 patients who were 75 years of age or older.

No overall differences in safety and efficacy were observed between patients and younger patients. In controlled clinical studies of pregabalin in fibromyalgia, 106 patients were 65 years of age or older. Although the adverse event profile was similar, the effect size for male reproductive organ hypotrophy was lower in older patients.

8.7 Renal Impairment
Pregabalin is eliminated primarily by renal excretion and dose adjustment is recommended for adult patients with renal impairment (see Dosage and Administration (2.7) and Clinical Pharmacology (12.3)). The use of pregabalin in pediatric patients with compromised renal function has not been studied.

9 DRUG ABUSE AND DEPENDENCE
9.1 Controlled Substances
Pregabalin is a Schedule I controlled substance.

Pregabalin is not known to be active at receptor sites associated with drugs of abuse. As with any CNS active drug, carefully evaluate patients for history of drug abuse and observe them for signs of pregabalin misuse or abuse (e.g., development of tolerance, dose escalation, drug-seeking behavior).


9.2 Abuse
In a study of recreational users (N=15) of sedative/hypnotic drugs, including alcohol, pregabalin (450 mg, single dose) received similar ratings of "good drug effect," "gratifying and pleasant" to a degree that was similar to diazepam (30 mg, single dose). Pregabalin was rated as "gratifying and pleasant" to a degree that was similar to diazepam (30 mg, single dose). In controlled clinical studies in 5500 patients, 4% of pregabalin-treated patients and 1% of placebo-treated patients overall reported euphoria as an adverse effect. In some patient populations studied, this reporting rate was higher and ranged from 1 to 12%.

9.3 Dependence
In controlled clinical studies, following abrupt or partial discontinuation of pregabalin, some patients reported symptoms including insomnia, nausea, headache or dizziness (see Warnings and Precautions (5.4)) consistent with physical dependence. In the postherpetic neuralgia study, in addition to these reported symptoms, other reported adverse reactions include, but are not limited to, seizures, depression, suicidal ideation and behavior, agitation, confusion, disorientation, psychotic symptoms, pain, swelling, tremor, dizziness, and vomiting.

10 OVERDOSE
Signs, Symptoms and Laboratory Findings of Acute Overdose in Humans
In the postherpetic neuralgia study, the most commonly reported adverse events observed with pregabalin when taken in overdose include decreased consciousness, depression/anxiety, confusion, dizziness, agitation, and decreased salivation. Signs and symptoms of overdose have been reported. Deaths have been reported in the setting of low pregabalin overdose and in combination with other CNS depressants.

Treatment of Management of Overdose
There is no specific antidote for overdose with pregabalin. If indicated, the administration of unabsorbed drug may be attempted with activated charcoal. In addition, the oral administration of calcium gluconate may be attempted if hypocalcemia is suspected. In addition, the oral administration of calcium gluconate may be attempted if hypocalcemia is suspected. In addition, the oral administration of calcium gluconate may be attempted if hypocalcemia is suspected.

11 DESCRIPTION
Pregabalin is chemically (S)-3-(aminomethyl)-5-methylhexanoic acid. The molecular formula is C10H19NO2 and the molecular weight is 159.23. The chemical structure of pregabalin is:



2500 mg/kg prior to and during mating with untreated females, a number of adverse reproductive and developmental effects were observed. These included decreased sperm counts and sperm motility, increased sperm abnormalities, reduced fertility, increased preimplantation embryo loss, decreased litter size, decreased fetal body weights, and an increased incidence of fetal abnormalities. Effects on sperm and fertility parameters were reversible in studies of this duration (3 to 4 months). The no-effect dose for reproductive toxicity in humans (100 mg/kg) was associated with a plasma pregabalin exposure (AUC) approximately 3 times human exposure at the maximum recommended dose (MRD) of 600 mg/day.

In addition, adverse reactions on reproductive organ (testes, epididymides) histopathology were observed in male rats exposed to pregabalin (500 mg/kg to 1250 mg/kg) in general toxicology studies of 4 weeks or greater duration. The no-effect dose for male reproductive organ hypotrophy was 250 mg/kg. The no-effect dose for male reproductive organ hypotrophy was 250 mg/kg. The no-effect dose for male reproductive organ hypotrophy was 250 mg/kg.

13.2 Animal Toxicology and/or Pharmacology
Skin lesions ranging from erythema to necrosis were seen in repeated-dose toxicology studies in both rats and monkeys. The etiology of these skin lesions is unknown. At the maximum recommended human dose (MRD) of 600 mg/day, there is a 2-fold safety margin for the dermatological lesions. The more severe dermatitis lesions involving necrosis were associated with plasma exposures (as expressed by plasma AUC) of approximately 3 to 8 times those achieved in humans given the MRD. No increase in incidence of skin lesions was observed in clinical studies.


Ocular Lesions
Ocular lesions characterized by retinal atrophy (including loss of photoreceptor cells) and/or corneal inflammation/vascularization were observed in two lifetime carcinogenicity studies in Wistar rats. These findings were observed in plasma pregabalin exposures (AUC) greater than or equal to 10 times those achieved in humans given the MRD. No increase in incidence of ocular lesions was observed in clinical studies.

14 CLINICAL STUDIES
14.1 Neuropathic Pain Associated with Diabetic Peripheral Neuropathy
The efficacy of the maximum recommended dose of pregabalin for the management of neuropathic pain associated with diabetic peripheral neuropathy was established in three double-blind, placebo-controlled, multicenter studies with three times a day dosing, two of which studied the maximum recommended dose. Patients were enrolled in either Type 1 or Type 2 diabetes mellitus and a diagnosis of diabetic neuropathy for 1 to 5 years.

A total of 89% of patients completed Studies DP1 and DP2. The patients had a minimum baseline pain score of greater than or equal to 4 on an 11-point numerical pain rating scale ranging from 0 (no pain) to 10 (worst possible pain). The baseline mean pain scores across the two studies ranged from 6.1 to 6.7. Patients were permitted up to 4 grams of acetaminophen per day as needed for pain. In addition to pregabalin, patients received placebo for pain.


Study DP1: This 8-week study compared pregabalin 25 mg, 100 mg, or 200 mg three times a day with placebo. Treatment with pregabalin 100 mg and 200 mg three times a day statistically significantly improved the endpoint mean pain score and increased the proportion of patients with at least a 50% reduction in pain score from baseline. There was no evidence of a greater effect on pain or quality of life from those achieved with placebo. A total of 63% of patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 1: Patients Achieving Various Levels of Improvement in Pain Intensity - Study DP1



Study DP2: This 8-week study compared pregabalin 100 mg three times a day with placebo. Treatment with pregabalin 100 mg three times a day statistically significantly improved the endpoint mean pain score and increased the proportion of patients with at least a 50% reduction in pain score from baseline. There was no evidence of a greater effect on pain or quality of life from those achieved with placebo. A total of 63% of patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.


Figure 2: Patients Achieving Various Levels of Improvement in Pain Intensity - Study DP2



14.2 Postherpetic Neuralgia
The efficacy of pregabalin for the management of postherpetic neuralgia was established in three double-blind, placebo-controlled, multicenter studies. These studies enrolled patients with neuropathic pain persisting for at least 3 months following healing of herpes zoster rash and a minimum baseline score of greater than or equal to 4 on an 11-point numerical pain rating scale ranging from 0 (no pain) to 10 (worst possible pain). Seventy-three percent of patients completed the studies. The baseline mean pain scores across the 3 studies ranged from 6.7 to 7.6. Patients were permitted up to 4 grams of acetaminophen per day as needed for pain. In addition to pregabalin, patients received placebo for pain.


Study PNH 1: This 13-week study compared pregabalin 75 mg, 150 mg, and 300 mg twice daily with placebo. Treatment with pregabalin 150 mg and 300 mg twice daily statistically significantly improved the endpoint mean pain score and increased the proportion of patients with at least a 50% reduction in pain score from baseline. There was no evidence of a greater effect on pain or quality of life from those achieved with placebo. A total of 63% of patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 3: Patients Achieving Various Levels of Improvement in Pain Intensity - Study PNH 1




Study PNH 2: This 8-week study compared pregabalin 100 mg or 200 mg three times a day with placebo. Treatment with pregabalin 100 mg and 200 mg three times a day statistically significantly improved the endpoint mean pain score and increased the proportion of patients with at least a 50% reduction in pain score from baseline. There was no evidence of a greater effect on pain or quality of life from those achieved with placebo. A total of 63% of patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 4: Patients Achieving Various Levels of Improvement in Pain Intensity - Study PNH 2



Study PNH 3: This 8-week study compared pregabalin 50 mg or 100 mg three times a day with placebo with doses assigned regardless of creatinine clearance. Treatment with pregabalin 50 mg and 100 mg three times a day statistically significantly improved the endpoint mean pain score and increased the proportion of patients with at least a 50% reduction in pain score from baseline. There was no evidence of a greater effect on pain or quality of life from those achieved with placebo. A total of 63% of patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 5: Patients Achieving Various Levels of Improvement in Pain Intensity - Study PNH 3



14.3 Adjunctive Therapy for Partial-Onset Seizures in Patients 1 Month of Age and Older
Adjunctive Therapy for Partial-Onset Seizures in Pediatric Patients 4 to Less Than 17 Years of Age
The efficacy of pregabalin as adjunctive therapy in partial-onset seizures was established in a 12-week, randomized, double-blind, placebo-controlled, multicenter study in pediatric patients 4 years to less than 17 years of age with partial-onset seizures with or without secondary generalization. During an 8-week baseline period, patients had to experience at least 6 partial-onset seizures with no seizure-free period exceeding 2 to 3 consecutive AEDs at baseline. Among the pregabalin-treated patients, 87% completed the double-blind phase of the study.

In this study, pregabalin 2.5 mg/kg/day (maximum 150 mg/day) and 10 mg/kg/day (maximum 600 mg/day) were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). Because of higher weight-normalized clearance in patients with body weight less than 30 kg (see Clinical Pharmacology (12.3)), the pregabalin dose was increased by 40% to 3.5 mg/kg/day for patients weighing less than 30 kg randomized to the 2.5 mg/kg/day group or to 14 mg/kg/day for patients randomized to the 10 mg/kg/day group.

Table 12: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Subsequent evaluations of the antiseizure efficacy of pregabalin showed no clinically important differences as a function of age, gender, or race. Adjunctive Therapy for Partial-Onset Seizures in Pediatric Patients 4 to Less Than 17 Years of Age
The efficacy of pregabalin as adjunctive therapy in partial-onset seizures was established in a 12-week, randomized, double-blind, placebo-controlled, multicenter study in pediatric patients 4 years to less than 17 years of age with partial-onset seizures with or without secondary generalization. During an 8-week baseline period, patients had to experience at least 6 partial-onset seizures with no seizure-free period exceeding 2 to 3 consecutive AEDs at baseline. Among the pregabalin-treated patients, 87% completed the double-blind phase of the study.

In this study, pregabalin 2.5 mg/kg/day (maximum 150 mg/day) and 10 mg/kg/day (maximum 600 mg/day) were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). Because of higher weight-normalized clearance in patients with body weight less than 30 kg (see Clinical Pharmacology (12.3)), the pregabalin dose was increased by 40% to 3.5 mg/kg/day for patients weighing less than 30 kg randomized to the 2.5 mg/kg/day group or to 14 mg/kg/day for patients randomized to the 10 mg/kg/day group.


Table 12: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: BD=twice daily; N=number. * 2.5 mg/kg/day: Maximum dose 150 mg/day. Includes patients less than 30 kg for whom dose was adjusted to 3.5 mg/kg/day. † 10 mg/kg/day: Maximum dose 600 mg/day. Includes patients less than 30 kg for whom dose was adjusted to 14 mg/kg/day.

There was evidence of a dose-response relationship for total daily doses of pregabalin (2.5 mg/kg/day and 10 mg/kg/day) compared with placebo. The 12-week study consisted of a 3-week dose adjustment phase and a 9-week dose maintenance phase. Treatment with pregabalin 10 mg/kg/day to 150 mg/day statistically significantly improved the endpoint weekly mean pain score, and increased the proportion of patients with at least a 50% reduction in pain score from baseline. The fraction of patients achieving various levels of improvement in pain intensity from baseline to Week 16 is presented in Figure 12. Some patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 11: Patients Achieving Various Levels of Improvement in Pain Intensity - Study S2 1




Study S2 2: This 16-week, randomized, double-blind, placebo-controlled, parallel-group, multicenter, flexible dose (150 mg/day to 600 mg/day), in increments of 150 mg study compared the efficacy, safety and tolerability of pregabalin compared with placebo. The 12-week study consisted of a 3-week dose adjustment phase and a 9-week dose maintenance phase. Treatment with pregabalin 10 mg/kg/day to 150 mg/day statistically significantly improved the endpoint weekly mean pain score, and increased the proportion of patients with at least a 50% reduction in pain score from baseline. The fraction of patients achieving various levels of improvement in pain intensity from baseline to Week 16 is presented in Figure 12. Some patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 12: Patients Achieving Various Levels of Improvement in Pain Intensity - Study S2 2



Figure 7: Seizure Reduction by Dose (All Partial-Onset Seizures) for Studies E1, E2, and E3



Subsequent evaluations of the antiseizure efficacy of pregabalin showed no clinically important differences as a function of age, gender, or race. Adjunctive Therapy for Partial-Onset Seizures in Pediatric Patients 4 to Less Than 17 Years of Age
The efficacy of pregabalin as adjunctive therapy in partial-onset seizures was established in a 12-week, randomized, double-blind, placebo-controlled, multicenter study in pediatric patients 4 years to less than 17 years of age with partial-onset seizures with or without secondary generalization. During an 8-week baseline period, patients had to experience at least 6 partial-onset seizures with no seizure-free period exceeding 2 to 3 consecutive AEDs at baseline. Among the pregabalin-treated patients, 87% completed the double-blind phase of the study.

In this study, pregabalin 2.5 mg/kg/day (maximum 150 mg/day) and 10 mg/kg/day (maximum 600 mg/day) were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). Because of higher weight-normalized clearance in patients with body weight less than 30 kg (see Clinical Pharmacology (12.3)), the pregabalin dose was increased by 40% to 3.5 mg/kg/day for patients weighing less than 30 kg randomized to the 2.5 mg/kg/day group or to 14 mg/kg/day for patients randomized to the 10 mg/kg/day group.


Table 12: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: BD=twice daily; N=number. * 2.5 mg/kg/day: Maximum dose 150 mg/day. Includes patients less than 30 kg for whom dose was adjusted to 3.5 mg/kg/day. † 10 mg/kg/day: Maximum dose 600 mg/day. Includes patients less than 30 kg for whom dose was adjusted to 14 mg/kg/day.


There was evidence of a dose-response relationship for total daily doses of pregabalin (2.5 mg/kg/day and 10 mg/kg/day) compared with placebo. The 12-week study consisted of a 3-week dose adjustment phase and a 9-week dose maintenance phase. Treatment with pregabalin 10 mg/kg/day to 150 mg/day statistically significantly improved the endpoint weekly mean pain score, and increased the proportion of patients with at least a 50% reduction in pain score from baseline. The fraction of patients achieving various levels of improvement in pain intensity from baseline to Week 16 is presented in Figure 12. Some patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 11: Patients Achieving Various Levels of Improvement in Pain Intensity - Study S2 1



Study S2 2: This 16-week, randomized, double-blind, placebo-controlled, parallel-group, multicenter, flexible dose (150 mg/day to 600 mg/day), in increments of 150 mg study compared the efficacy, safety and tolerability of pregabalin compared with placebo. The 12-week study consisted of a 3-week dose adjustment phase and a 9-week dose maintenance phase. Treatment with pregabalin 10 mg/kg/day to 150 mg/day statistically significantly improved the endpoint weekly mean pain score, and increased the proportion of patients with at least a 50% reduction in pain score from baseline. The fraction of patients achieving various levels of improvement in pain intensity from baseline to Week 16 is presented in Figure 12. Some patients experienced a decrease in pain as early as Week 1, which persisted throughout the study.

Figure 12: Patients Achieving Various Levels of Improvement in Pain Intensity - Study S2 2



14.4 Management of Fibromyalgia
The efficacy of pregabalin for the management of fibromyalgia was established in a 14-week, double-blind, placebo-controlled, multicenter study (F1) and one six-month, randomized withdrawal study (F2). Studies F1 and F2 enrolled patients with a diagnosis of fibromyalgia using the American College of Rheumatology (ACR) criteria (history of widespread pain for 3 months, and pain present at 11 or more of the 18 tender point sites). The study in Study F1 was a 14-week, double-blind, placebo-controlled study. The study in Study F2 was a 6-month, randomized withdrawal study.

In this study, pregabalin 75 mg/kg/day and 150 mg/kg/day were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). The 14-week study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase.

Table 13: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: N=number of patients. A significant improvement in partial-onset seizure rate was observed for pregabalin 14 mg/kg/day group compared with placebo. Patients treated with pregabalin 7 mg/kg/day did not show improvement relative to placebo.

Responder rates (>50% or greater reduction in partial-onset seizure frequency) were a secondary efficacy parameter; patients treated with pregabalin 14 mg/kg/day showed numerical improvement compared with placebo. The responder rates for patients treated with pregabalin 7 mg/kg/day and 14 mg/kg/day were 53.6%, 30.5%, and 41.5%, for pregabalin 14 mg/kg/day, Pregabalin 7 mg/kg/day, and placebo, respectively.

The efficacy of pregabalin for management of fibromyalgia was established in a 14-week, double-blind, placebo-controlled, multicenter study (F1) and one six-month, randomized withdrawal study (F2). Studies F1 and F2 enrolled patients with a diagnosis of fibromyalgia using the American College of Rheumatology (ACR) criteria (history of widespread pain for 3 months, and pain present at 11 or more of the 18 tender point sites). The study in Study F1 was a 14-week, double-blind, placebo-controlled study. The study in Study F2 was a 6-month, randomized withdrawal study.

In this study, pregabalin 75 mg/kg/day and 150 mg/kg/day were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). The 14-week study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase.

Table 13: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: N=number of patients. A significant improvement in partial-onset seizure rate was observed for pregabalin 14 mg/kg/day group compared with placebo. Patients treated with pregabalin 7 mg/kg/day did not show improvement relative to placebo.

Responder rates (>50% or greater reduction in partial-onset seizure frequency) were a secondary efficacy parameter; patients treated with pregabalin 14 mg/kg/day showed numerical improvement compared with placebo. The responder rates for patients treated with pregabalin 7 mg/kg/day and 14 mg/kg/day were 53.6%, 30.5%, and 41.5%, for pregabalin 14 mg/kg/day, Pregabalin 7 mg/kg/day, and placebo, respectively.

The efficacy of pregabalin for management of fibromyalgia was established in a 14-week, double-blind, placebo-controlled, multicenter study (F1) and one six-month, randomized withdrawal study (F2). Studies F1 and F2 enrolled patients with a diagnosis of fibromyalgia using the American College of Rheumatology (ACR) criteria (history of widespread pain for 3 months, and pain present at 11 or more of the 18 tender point sites). The study in Study F1 was a 14-week, double-blind, placebo-controlled study. The study in Study F2 was a 6-month, randomized withdrawal study.

In this study, pregabalin 75 mg/kg/day and 150 mg/kg/day were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). The 14-week study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase.

Table 13: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: N=number of patients. A significant improvement in partial-onset seizure rate was observed for pregabalin 14 mg/kg/day group compared with placebo. Patients treated with pregabalin 7 mg/kg/day did not show improvement relative to placebo.

Responder rates (>50% or greater reduction in partial-onset seizure frequency) were a secondary efficacy parameter; patients treated with pregabalin 14 mg/kg/day showed numerical improvement compared with placebo. The responder rates for patients treated with pregabalin 7 mg/kg/day and 14 mg/kg/day were 53.6%, 30.5%, and 41.5%, for pregabalin 14 mg/kg/day, Pregabalin 7 mg/kg/day, and placebo, respectively.

The efficacy of pregabalin for management of fibromyalgia was established in a 14-week, double-blind, placebo-controlled, multicenter study (F1) and one six-month, randomized withdrawal study (F2). Studies F1 and F2 enrolled patients with a diagnosis of fibromyalgia using the American College of Rheumatology (ACR) criteria (history of widespread pain for 3 months, and pain present at 11 or more of the 18 tender point sites). The study in Study F1 was a 14-week, double-blind, placebo-controlled study. The study in Study F2 was a 6-month, randomized withdrawal study.

In this study, pregabalin 75 mg/kg/day and 150 mg/kg/day were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). The 14-week study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase.

Table 13: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

Abbreviations: N=number of patients. A significant improvement in partial-onset seizure rate was observed for pregabalin 14 mg/kg/day group compared with placebo. Patients treated with pregabalin 7 mg/kg/day did not show improvement relative to placebo.

Responder rates (>50% or greater reduction in partial-onset seizure frequency) were a secondary efficacy parameter; patients treated with pregabalin 14 mg/kg/day showed numerical improvement compared with placebo. The responder rates for patients treated with pregabalin 7 mg/kg/day and 14 mg/kg/day were 53.6%, 30.5%, and 41.5%, for pregabalin 14 mg/kg/day, Pregabalin 7 mg/kg/day, and placebo, respectively.

The efficacy of pregabalin for management of fibromyalgia was established in a 14-week, double-blind, placebo-controlled, multicenter study (F1) and one six-month, randomized withdrawal study (F2). Studies F1 and F2 enrolled patients with a diagnosis of fibromyalgia using the American College of Rheumatology (ACR) criteria (history of widespread pain for 3 months, and pain present at 11 or more of the 18 tender point sites). The study in Study F1 was a 14-week, double-blind, placebo-controlled study. The study in Study F2 was a 6-month, randomized withdrawal study.

In this study, pregabalin 75 mg/kg/day and 150 mg/kg/day were compared to placebo. Administration of each daily dose was divided into two equal doses (twice a day dosing). The 14-week study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase. The 6-month study consisted of a 24-hour run-in period followed by the double-blind phase.

Table 13: Seizure Response in Controlled Adjunctive Partial-Onset Seizure Study in Pediatric Patients 4 to Less Than 17 Years of Age

Table with 6 columns: Daily Dose of Pregabalin, N, Median Baseline Seizure Frequency/24 hrs, Median % Change from Baseline, % Difference Relative to Placebo, p-value, versus placebo. Rows include Placebo, 2.5 mg/kg/day (BD), and 10 mg/kg/day (BD).

16 HOW SUPPLIED/STORAGE AND HANDLING
16.1 HOW SUPPLIED/STORAGE AND HANDLING
White to off white powder filled in size "4" hard gelatin capsules with white opaque colored cap and white opaque colored body imprinted "SC" on cap and "359" on body with black ink, available in:

Bottles of 30: NDC 50228-353-30
Bottles of 90: NDC 50228-353-90
Bottles of 100: NDC 50228-353-100
Bottles of 300: NDC 50228-353-300

50 mg capsules: White to off white powder filled in size "3" hard gelatin capsules with white opaque colored cap and white opaque colored body imprinted "SC" on cap and "352" on body with black ink, available in:

Bottles of 30: NDC 50228-352-30
Bottles of 90: NDC 50228-352-90
Bottles of 100: NDC 50228-352-100
Bottles of 300: NDC 50228-352-300

100 mg capsules: White to off white powder filled in size "1" hard gelatin capsules with orange opaque colored cap and white opaque colored body imprinted "SC" on cap and "354" on body with black ink, available in:

Bottles of 30: NDC 502